




(51) International Patent Classification ³ : A61K 39/29; C12N 7/00, 7/02 G01N 33/54	A1	(11) International Publication Number: WO 84/ 01107
		(43) International Publication Date: 29 March 1984 (29.03.84)
<p>(21) International Application Number: PCT/US83/01412</p> <p>(22) International Filing Date: 16 September 1983 (16.09.83)</p> <p>(31) Priority Application Number: 418,708</p> <p>(32) Priority Date: 16 September 1982 (16.09.82)</p> <p>(33) Priority Country: US</p> <p>(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; Fruit Street, Boston, MA 02114 (US).</p> <p>(72) Inventors: WANDS, Jack ; 210 Varick Road, Waban, MA 02168 (US). SHAFRITZ, David, A. ; 4 Pheasant Run, Larchmont, NY 10538 (US).</p> <p>(74) Agents: GOLDSTEIN, Jorge, A. et al.; Oblon, Fisher, Spivak, McClelland & Maier, Crystal Square Five, Suite 400, 1755 South Jefferson Davis Highway, Arlington, VA 22202 (US).</p>		<p>(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), JP, LU (European patent), NL (European patent), SE (European patent).</p> <p>Published <i>With international search report.</i></p>



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	KR	Republic of Korea
AU	Australia	LI	Liechtenstein
BE	Belgium	LK	Sri Lanka
BG	Bulgaria	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Republic	MG	Madagascar
CG	Congo	MR	Mauritania
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SD	Sudan
FR	France	SE	Sweden
GA	Gabon	SN	Senegal
GB	United Kingdom	SU	Soviet Union
HU	Hungary	TD	Chad
JP	Japan	TG	Togo
KP	Democratic People's Republic of Korea	US	United States of America

-1-

DescriptionNon A, Non B Hepatitis Virus, Methods of
Identification, Purification, Characterization,
Diagnosis and Immunization5 Technical Field

Part of the present invention was developed with funds obtained from the following sources: AG-04145, AA-02666, AM-17702, AM 17609, CA-32605, AM-07218, 1-K02-AA-00048, from the United States
10 Government.

The present invention deals with the identification, isolation, characterization, purification and use of non A, non B hepatitis virus, as well as diagnostic methods and vaccines methods therefor.

15 Background Art

The name non A, non B hepatitis is given to acute and chronic cases of viral hepatitis in humans which occur in the absence of infection with any known or serologically identifiable virus associated with
20 hepatitis B (HBV) or hepatitis A (HAV). The characteristics of non-A, non-B (hereinafter "NANB") hepatitis are well described in Dienstag et al, Chapter 302 of Harrison's "Principles of Internal Medicine", 9th Ed, McGraw-Hill Book Co., 1980, pp. 1459-1467, and
25 by Robinson, W. S., "The Enigma of Non-A, Non-B Hepatitis", The Journal of Infect. Dis., Vol. 145 No.



-2-

3, pp. 387-395 (1982). These two articles are herein incorporated by reference, and the following comments are extracted therefrom.

5 Sensitive serologic tests for identifying both types A and B hepatitis have led to the identification of hepatitis cases with incubation periods and modes of transmission consistent with an infectious disease, but without serologic evidence of hepatitis A or B infection.

10 Transmission of the disease to chimpanzees has clearly established that many of the cases are caused by one or more infectious agents. There have been intensive efforts in many laboratories throughout the world in the past few years to identify and
15 characterize agents that are responsible for these infections.

Clinical diagnosis of NANB hepatitis is made by excluding infection with known hepatitis viruses and other known factors that cause hepatitis. The
20 infection occurs with high frequency after blood transfusion or parental drug abuse, in person to person contact and in other settings that are also associated with HBV infections. Endemic and apparently epidemic disease has also been observed without obvious overt
25 parental transmission.

Despite these advances and intensive efforts to date, no etiologic agent of NANB hepatitis has been unequivocally identified as an antigenic ultra-structural or molecular entity. This result suggests
30 that the concentration of viral antigen in the serum of patients with NANB hepatitis may be much lower than



that of HBV antigen in patients with hepatitis B, or that appropriate reagents or methods have not been heretofore described to identify the virus, its proteins, or its genetic material.

5 The most important experimental advance in this field to date has been the transmission of NANB hepatitis agents to chimpanzees. This provided a direct demonstration of a transmissible agent, associated with NANB hepatitis, in an animal model of
10 the disease (See, for example, Alter, H.J. et al, Lancet 1: 459-463 (1978), Tabor, E. et al, ibid 1: 463-466 (1978), Hollinger, F.B. et al, Intervirology 10: 60-68 (1978), or Bradley D.W. et al, J. Med. Virol. 3: 253-269 (1979), all of which are herein incorporated by
15 reference).

Despite the fact that NANB hepatitis has been transmitted to experimental animals, no virus or other infectious agent(s) has been physically identified with certainty prior to this invention. Although detection
20 of apparently unique antigen/antibodies systems in the sera of patients and chimpanzees with NANB hepatitis have been reported, the results have been difficult to confirm, and none of these tests has clearly identified sera known to contain NANB agents (see for example,
25 Vitvitski, L. et al, Lancet 22: 1263-1267 (1979), Kabiri, M. et al, Lancet 2: 221-224 (1979), Tabor E., J. Med. Virol. 4: 161-169 (1979) and Chircu, L.V. et al, J. Med. Virol. 6: 147-151 (1980).) In addition to antigen, virus-like particulate structures have been
30 observed by electron microscopy in serum and liver of humans and chimpanzees infected with NANB hepatitis (see for example Bradley, D.W., J. Med. Virol. 3: 253-269 (1979) and Bradley, D.W. et al, J. Med. Virol. 6:



85-201 (1980)).

An evaluation of all of these studies has been made by Robinson, supra in J. Inf. Dis. Vol. 145, (1982) who stated that: "Without more definitive
5 evidence concerning these particles and because numerous investigators have failed to confirm these findings it is not possible at this time to conclude that any HBV-like virus is ever a cause of NANB hepatitis."

10 In view of all of the above, it is quite clear that there exists at present a great need to identify, isolate and characterize the etiologic agent(s) causative of NANB hepatitis. A need also exists for
15 accurate and unambiguous identification and detection techniques therefor, which will help in the quick and accurate diagnosis of the disease.

Disclosure of the Invention

It is therefore an object of the invention to provide an accurate and specific characterization of
20 the etiologic agent of NANB hepatitis.

It is another object of the invention to provide for a method of identifying and detecting the etiologic agent of NANB hepatitis in samples.

Still another object of the invention is to
25 provide for a method of diagnosing NANB hepatitis in animals.

Yet another object of the invention is to provide a vaccine against NANB hepatitis, and a method



of immunization which comprises the use of such vaccine.

Still another object of the invention is to provide a method for the purification of NANB hepatitis
5 virus.

These and other objects of the invention as will hereinafter become more readily apparent have been attained by providing:

An attenuated or inactivated form of a DNA
10 virus which, in unattenuated or inactivated form has the following characteristics:

- 1)molecular weight greater than 2×10^6 daltons;
- 15 2)substantial immunoreactivity towards commercially available anti HBsAg IgM monoclonal antibody 5D3 (obtained from a RadioImmunoAssay Test Kit Sold by Centocor, USA; see below);
- 20 3)substantially no immunoreactivity towards an anti HBsAg monoclonal antibody obtained from cell line ATCC CRL 8018;
- 4)concentration dependent binding capacity towards polyclonal IgG anti-HBsAg antibodies, which increases with increased
25 concentration of said DNA virus;
- 5)discrete particulate form when observed by immunoelectron microscopy in the presence of 5D3 IgM antibodies;
- 30 6)a polypeptide profile on sodium dodecyl sulfate polyacrylamide gels, when affinity purified with IgM antibody from cell line ATCC HB 8058, comprising bands at about



-6-

50,000, about 23,000 and about less than 20,000 molecular weight;

7)the DNA of said virus showing partial sequence homology with hepatitis B virus DNA by molecular hybridization; and
8)said DNA virus showing, in chimpanzees, infectivity having the characteristics of non A, non B hepatitis.

Another object of the invention has been attained by providing a method of detecting the presence of non A, non B hepatitis virus in the sample of an animal which comprises A) confirming the presence of said virus in said sample, and B) distinguishing said virus from hepatitis B virus.

Another object of the invention has been obtained by providing a method of purifying NANB virus from an animal sample by immunoaffinity chromatography wherein the immunosorbent antibody is a monoclonal antibody having substantial immunoreactivity towards said NANB virus.

The present invention also provides vaccines and vaccination methods utilizing live, attenuated or inactivated forms of the NANB virus.

Brief Description of the Drawings

The present invention will become better understood by reference to the accompanying description when interpreted in view of the following drawings, wherein:

FIGURE 1 is a representative example of binding activity isolated from human serum after elution of the



-7-

monoclonal 5D3-IgM anti-HBs affinity column with glycine-HCl buffer (pH 2.6). Symbols are as follows: (●) CPM bound in the eluate by monoclonal 5D3-5D3 radioimmunoassay (RIA); (▲) CPM bound in the monoclonal assay on fractionated serum passed through the affinity column and eluted with Pi/NaCl; (■) binding profile of glycine. HCl eluate when analyzed by a commercial radioimmunoassay kit (polyclonal antibodies, AUSRIA II), See Example 1;

FIGURE 2 shows the binding profiles exhibited by five monoclonal RIAs in human serum derived from three patients (See Example 1). All monoclonal RIA antibodies are reactive with serial dilutions of serum and indicate that such IgG and IgM monoclonal anti-HBs recognize determinants present on HBsAg (patient D) (Right). In contrast, only the 5D3-5D3 monoclonal assay shows high binding values in serial dilutions of serum from patients B and C (left). Description and characterization of the monoclonal antibodies is given in the text, infra;

FIGURE 3 shows the polypeptide profile on sodium dodecyl sulfate/polyacrylamide gel of the affinity purified material from patients' serum (Example 1): Patient A, acute hepatitis; Patient B, chronic active hepatitis; Patient C, blood donor; and Patient D, HBsAg-positive chronic acute hepatitis. There are three similar polypeptides in all four specimens (noted as 1, 2 and 3). Polypeptide 1 has a M_r of 50,000 and polypeptides 2 and 3 have M_r s of 22,000-23,000. In Sample D there is a polypeptide of M_r 27,000-30,000. However, in Samples A, B, and C there are three additional majors protein bands not observed in Sample D; one has a M_r of approximately



-8-

80,000 (compared with 5D3 heavy chain) and of the two others, the first has a M_r slightly greater than 23,000 and the second a M_r less than 20,000. The heavy and light chains of the 5D3 antibody do not comigrate with
5 any other proteins from Patients A, B, C and D.

FIGURE 4 shows the comparative inhibition of binding to HBsAg determinants by monoclonal anti-HBs antibodies (See Example 2). The IgG anti-HBs antibodies 2C6 and 5C3 have no effect on the binding of
10 5D3 to HBsAg-related determinant(s), whereas 3D4, an IgM anti-HBs, partially inhibited 5D3 binding;

FIGURE 5 shows a comparison of the IgM monoclonal RIA (■) with polyclonal antibody AUSRIA II (▲) in a patient with acute hepatitis B and immune
15 complex disease. ---O---SGOT (serum alanine amino transferase). S/N, signal -to-noise ratio defined as CPM bound in experimental samples/CPM bound in controls (See Example 3);

FIGURE 6 shows the clinical and virologic
20 course of non-A, non-B hepatitis in a chimpanzee. The elevations of ALT precede the appearance of antigen and HBV-related DNA sequences in the blood (See Example 4, for this and for FIGURES 7-10);

FIGURE 7 shows the detection of hepatitis B-
25 virus-DNA related sequences by molecular hybridization analysis in 250 μ L serum from two chimpanzees with non-A, non-B hepatitis. *: Denotes positive results with a recombinant, cloned HBV-DNA probe. Spots 1, 2 and 3 were negative for HBV-DNA but positive for antigen by
30 IgM anti-HBs radioimmunoassays (see Figures 9 and 10). Spots 4, 5 and 6 were positive for both antigen



and HBV-DNA related sequences in serum on days 47 , 58 and 64 (Figure 6). Spots 8 and 9 were also positive for antigen and HBV-related sequences on days 190 and 204 in the second animal (Figures 8). Spots 7 and 10 are negative controls;


FIGURE 8 shows the clinical and virologic course of non A, non B hepatitis in a chimpanzee. Despite infection, as shown by the presence of viral antigen and HBV-DNA related sequences in serum, ALT values remained normal;

FIGURE 9 shows the clinical and virologic course of non A, non B hepatitis in a chimpanzee. Similar to Figure 6, ALT elevations precede the appearance of antigen detected by the monoclonal IgM anti-HBs radioimmunoassays by approximately 45 days. HBV-DNA related sequences were not detected in this animal;

FIGURE 10 shows the clinical and virologic course of non A, non B hepatitis in a chimpanzee. The appearance of three well defined peaks of antigenemia as measured by the monoclonal IgM anti-HBs radioimmunoassays should be noted.

Best Mode for Carrying Out the Invention

The present invention is based on the discovery of highly specific and accurate tests for the identification and characterization of the causative agent of non A, non B hepatitis. The inventors have made use of a variety of analytical techniques to characterize NANB hepatitis virus and distinguish the same from hepatitis virus A (HVA) and hepatitis virus B



-10-

(HBV). These techniques include physical-chemical properties, immunological properties, genetic characterization and infectivity characterization.

The discovery of NANB hepatitis virus was made by detecting its presence in the blood of persons with the clinical signs of hepatitis but no serologic identification by any of the prior art immunoassay techniques using polyvalent IgG antibodies. A series of monoclonal antibody screening tests were then developed with alternatively positive and negative binding for various different monoclonal antibodies, which can readily characterize and detect NANB virus and distinguish the same from hepatitis B virus.

In the discussions that follow, mention is made to a number of antibodies, both monoclonal and polyvalent. For clarity purposes and reference, the following summarizes the nature, origin and type of these antibodies:

1) 5D3: Represents a monoclonal IgM antibody against HBsAg, obtained commercially as polystyrene bead bound IgM antibody, present in the "RIA Test Kit for Detection of HBsAg, (US lic No. 889)", sold by Centocor, 244 Great Valley Parkway, Malvern, PA USA 19355. Reference is made to this antibody in U.S. Patent 4,271,145 to Wands et al, as well as in Wands et al, Proc. Nat. Acad. Sci., USA Vol. 78: 1214-1218, February 1981, both of which are herein incorporated by reference.

2) 3D4: Represents a monoclonal IgM antibody having specificity against HBsAg (i.e. anti-HBsAg), obtained from cell line 3D4 on deposit at the ATCC with deposit number HB-8170.



-11-

3) 1F8: Represents a monoclonal IgM anti-HBsAg antibody derived from cell line 1F8 on deposit at the ATCC have deposit number CRL 8018. This antibody is described in the aforementioned Wands et al U.S. Patent 4,271,145 and Wands et al PNAS Vol. 78, February 1981 paper.


4) 5C11: Represents a monclonal IgG₁ anti-HBsAg antibody obtained from cell line 5C11 deposited at the ATCC with deposit number HB-8171.

10 5) AUSRIA II: Represents antibodies from a commercially available HBsAg test kit (Abbott) containing polyclonal IgG antibodies.

These and other other mentioned antibodies are also described in PCT patent publication: Serial No. PCT/US 81/01270, filed September 19, 1980 to Wands, Zurawski and Schoemaker, for "Immunoassay Utilizing Monoclonal High Affinity IgM Antibodies", published April 1, 1982 (No WO 82/01072).

Also, a bacterial culture containing a recombinant plasmid with HBV-DNA sequences, pAO1 HBV is on deposit with ATCC Number 31873.

The NANB virus can be isolated from either a human or other animal host, e.g., chimpanzee, marmoset, and other suitable hosts for NANB virus, which is infected with NANB hepatitis. The presence of the NANB virus has been implicated in the prior art by excluding identifiable hepatitis viruses (HAV, HBV, Epstein-Barr virus, cytomegalovirus and others) and other etiologic factors (for example, hepatotoxic drugs and chemicals). Exclusion of other viruses mentioned above can still be used to suggest but not establish the



-12-

presence of the NANB infectious agent in the host. However, with the advent by the present invention of highly specific tests for NANB virus it is preferred to
5 utilize these, see infra.

Preferably, affinity chromatography using monoclonal 5D3 IgM anti-HBs can be utilized for the purification and subsequent characterization of the antigen composition. A suitable material is obtained
10 by coupling 5D3 to Sepharose 6B®. Serum from an appropriate host is placed in contact with the monoclonal antibody on the solid phase support and the material is incubated for several hours at room temperature. The supports are then extensively washed
15 with an appropriate physiological buffer (e.g., PBS-phosphate buffer saline) at a physiological pH. Column fractions can then be collected with an acidic buffer (for example pH 2-3). The pH of each fraction is adjusted to physiological pH, and binding activity is
20 determined with the appropriate antibodies. Peak fractions exhibiting the highest binding activity can then be pooled to collect the NANB virus. Virus can also be isolated from the supernatant of any cell culture (e.g., bacteria, yeasts and other eukarotic
25 cells infected with said virus or viral DNA) or fermentation broth producing the same.

The NANB virus can be characterized and identified by at least four different characteristics, each of which is described in turn hereinbelow.

30 Physical-chemical characteristics. NANB virus particle has a molecular weight of approximately 2×10^6 as determined by Sepharose 4B® chromatography. The virus appears as distinct particles by immunoelectron



-13-

microscopy. When the virus is isolated from serum by the 5D3-IgM affinity chromatography described above, prominent spiculated particles at 220,000 X are observed, suggesting the presence of 5D3 anti-HBs on their surface. When affinity purified material is applied to sodium dodecyl sulfate/10% polyacrylamide gels and compared to HBsAg virus it is seen that there are similar polypeptides in all specimens at a molecular weight of about 50,000 and at molecular weights of 22,000-23,000. In addition, however, NANB virus shows three additional major protein bands not observed in HBsAg, one has a Mr of approximately 80,000 and, of the two others, the first has a molecular weight slightly greater than 23,000 and the second has a molecular weight less than 20,000. (See Figure 3).

Immunological characteristics. NANB virus reacts with some monoclonal antibodies having specificity, i.e., immunoreactivity, for distinct HBsAg-related epitopes, and not with other such anti-HBsAg monoclonal antibodies. For example, NANB will cross react at all concentrations with antibody 5D3 or with antibody 3D4 (both of which are monoclonal IgM anti-HBsAg antibodies). On the other hand, NANB will not cross-react with antibody 1F8 (also a monoclonal IgM having specificity against HBsAg) or with monoclonal 5C11 (a IgG₁ antibody). This serves to clearly distinguish NANB virus from hepatitis B virus, which reacts with these monoclonal antibodies. Immunoreactivity of NANB with polyvalent anti-HBs antibodies (commercially available AUSRIA II) is concentration dependent. At concentrations of about 1 ng to 100 ng, the polyvalent IgG antibodies do not detect or bind the NANB virus. Upon concentration of NANB by about 100 fold or larger than these, binding



-14-

and detection by polyvalent IgG can be observed. However, in some instances, polyvalent anti-HBs antibodies do not detect or bind to NANB hepatitis serum even after enrichment by affinity chromatography and 100 fold concentration as described above. Preincubation of NANB with 5D3 anti-HBs at these higher concentrations blocks the binding by conventional polyvalent anti-HBs.

Genetic Characteristics.

10 The DNA sequence of NANB virus is partially homologous related but not identical to HBV-DNA. It can thus be detected by hybridization with a purified HBV-DNA probe. See, infra.

Infectivity Characteristics. NANB virus having
15 the above physico-chemical, morphological, immunologic and genetic characteristics is infectious. Infectivity studies of viral hepatitis are positive in chimpanzees and in man. The characteristics for the infection are different than those normally seen for HBV or HAV. The
20 incubation period, as defined from inoculation of infectious material to the appearance of virus or viral protein in the blood, is longer than previously recognized. Alanine aminotransferase (ALT) elevation precedes the appearance of antigenemia by several
25 weeks. Antigenemia may occur in the absence of ALT elevations, a phenomenon observed in man. A chronic viral carrier state in man and chimpanzees may occur. The period of antigenemia and/or viremia appears to persist for weeks to months and usually disappears with
30 recovery. Antigenemia is still detectable in the resolution phase of illness when ALT levels are normal, a similarity to HBV infection in man. Several episodes of antigenemia may occur during the course of infection. Pre-existing anti-HBs is not protective in



-15-

the animal, confirming that NANB virus is sufficiently different in antigenic composition from HBV. (See Figures 6, and 8-10).

5 It should be noted, of course, that the
aforementioned characteristics are but only one
possible set. Obviously, as more of these
characteristics are researched and discovered it may be
possible to characterize the virus, for example, by
additional monoclonal antibodies, or DNA probes
10 including one or more that do not cross react at all
with hepatitis B. This possibility, however, is fully
contemplated in the present application which, when
pertaining to the virus per se, is meant to cover the
virus itself regardless of any additional or even novel
15 identifying tests.

The NANB virus characteristics can be used to
develop highly sensitive and accurate tests for
detecting the presence of NANB virus in animal samples,
such as blood - especially blood to be transfused-,
20 serum, urine, milk, tissue samples, feces, and the
like. Particularly useful is the detection of NANB
virus in animal serum, especially human serum, and
products derived from human blood, such as red blood
cells, plasma, platelet concentrates, clotting factor
25 concentrates and the like, for the diagnosis of NANB
hepatitis. Also particularly useful is the detection
of NANB virus in samples of blood from blood donors, to
screen for the possibility of transmission of NANB
hepatitis infection to recipients.

30 The availability of purified NANB virus allows
for the development of immunoassay procedures. The
appropriate antibodies can be used in any of the



-16-

multiple immunoassay procedures currently available to the art (see for example, T. Chard "An Introduction to Radioimmunoassay and Related Techniques", North-Holland 1978, or Schuurs, A.H.W.M, et al, "Enzyme Immunoassay", Clin. Chim. Acta 81: 1-40 (1977), both of which are herein incorporated by reference). For example, the presence of the virus in a sample can be detected by radioimmunoassay, enzyme immunoassay, or latex agglutination immunoassay. The technique utilized can be competitive, "sandwich" (forward, reverse or simultaneous), double antibody, or enzyme cascade, all of which are well known to those of skill in the art. It may be useful for certain techniques to prepare, by art known methods, detectably labeled NANB virus such as NANB labeled with a radiolabel (I^{125} , C^{14} , H^3 , P^{32} , etc.), with an enzyme (alkaline phosphatase, peroxidase, etc.) with a fluorescent probe, and the like. The antibodies can be either in solution or immobilized, such as for example, on the inside of tubes, on polymer or glass beads, on plastic strips, and the like.

Detection can also be carried out by hybridization analysis using a detectably labeled probe. The genetic information or code of a specific virus comprises a nucleic acid which may be composed of a polymer of ribonucleotides (RNA) or deoxyribonucleotides (DNA). It is known that nucleotide molecules that are complementary to one another can interact in solution by "hydrogen-bonding" to form stable base pairs. Thus, adenine recognizes thymidine and guanine recognizes cytosine. When two single-stranded, complementary, DNA molecules are present in a solution under conditions in which the complementary nucleotides can recognize one another,



-17-

these molecules will interact to form a stable duplex structure. This duplex is resistant to attack by certain nucleases which totally degrade single-stranded DNA. It is therefore possible to ascertain with great
5 precision the extent of duplex formation. This interaction of base sequences in polynucleotides reacting in solution is referred to as "reannealing" or "molecular hybridization" and can be performed under specific and sensitive conditions in which false
10 interactions do not occur.


For substantially stable and recognizable hybrids to be formed, minimum complementary sequence lengths of approximately 50-100 nucleotides or more often 100-200 nucleotides are required. The ability to
15 form such hybrids appears to depend on the experimental conditions of the hybridization reaction (ionic strength, polarity, pH and temperature of the hybridization solution), the concentration of the complementary nucleic acid molecules and the length of
20 time of the incubation. Another variable in the reaction is the physical state of the DNA in the test sample, in that it can be in solution or fixed to a solid support matrix such as a nitrocellulose filter paper. In the latter case, the rate of hybridization
25 between the detecting probe and the test sample of DNA affixed to the solid support surface is slowed by approximately 30%. The latter method is, however, extremely sensitive for detection of hybridizing sequences and with a $[^{32}\text{P}]$ radioactively labeled DNA
30 probe of specific activity $2-4 \times 10^8$ cpm per μg DNA, as easily obtained by workers skilled in the art, a 2-5 mm diameter circular spot on a nitrocellulose filter containing 0.1 μg (10^{-13} gm) of specific DNA sequence or less can be detected.



-18-

Depending on the various factors mentioned above, the hybridization reaction can be performed under very stringent conditions, so that a perfect or near perfect match in complementary DNA sequence is required or under less stringent conditions in which only a partial match is required. As the conditions for stringency of hybridization are relaxed, nucleic acid molecules of lesser and lesser sequence homology will form hybrids. This, of course, decreases the specificity of the reaction and raises the chances of false positive results. Therefore, in the preferred embodiment, hybridization conditions of high stringency have been used, so that only molecules with sequence regions of approximately 100-200 nucleotides or more in common with or nearly identical to HBV-DNA will form stable and detectable hybrids on a nitrocellulose filters. This enables the use of the hybridization method to identify DNA molecules in any cell, tissue, tissue extract, serum, plasma, body fluid, secretum, semen, breast milk, vaccine or the like, containing DNA molecules or genetic information closely related, nearly identical or identical to NANB-DNA.

As disclosed in this invention, NANB hepatitis virus(es) contain sequences closely related to HBV-DNA and can be detected by hybridization with a purified and suitably labeled HBV-DNA probe. DNA or RNA molecules which are not closely related to HBV-DNA will not be identified or detected by this method. These methods, considerations and conditions as well as many variations in hybridization technology as well as means to detect, isolate and identify hybrids are well known to those skilled in the art. Details concerning the preparation of the recombinant HBV-DNA probe, the labeling of the probe, the hybridization conditions are



-19-

described in Chakraborty et al, Nature, Volume 286, No. 5772, pages 531-533, July 31, 1980; Shouval et al, Proceedings National Academy of Science (PNAS) U.S.A., Volume 77, No. 10, pages 6147-6151, October, 1980; 5 Shafritz and Kew, Hepatology, Volume 1, No. 1, pages 1-8, Jan.-Feb., 1981; Shafritz, D.A. et al, New Engl. J. Med., 305:1067-1073, 1981; and copending U.S. patent application serial number 249,369, filed March 31, 1981 entitled Diagnostic Test for Hepatitis B Virus; all 10 hereby incorporated by reference.

Regardless of the technique(s) used, the detection of the virus in a sample is carried out by an overall two step test, which not only serves to confirm its presence but also distinguishes it from HBV, with 15 which it is closely related.

For example, the detection test can comprise a first step of testing for immunoreactivity with an antibody such as 5D3 or 3D4, with which NANB virus is reactive, followed by a second step of immunoassay with 20 an antibody such as 5C11 or 1F8 with which NANB virus is not cross reactive, but HBV is.

Another two step test comprises a first immunoassay step using an antibody such as 5C11 or 1F8 (showing no cross reactivity), followed by DNA 25 hybridization using an HBV-DNA or an NANB-DNA (see infra) detectably labeled probe (e.g., ³²P or biotin-labeled probe).

An alternative test is a two step methodology wherein the first step is an immunoassay with 5D3 or 30 3D4 monoclonal IgM, followed by studying the infectivity characteristics in chimpanzees.



-20-

Alternatively, a two step analysis can be used with the first step being an immunoassay with 5D3 and in a second step a polyacrylamide gel on sodium dodecyl sulfate seeking the differential proteins present in
5 NANB and not present HBsAg.

There are obviously other possibilities, such as procedures utilizing more than two steps, for example, screening with 5D3, 3D4, 5C11, 1F8, testing for hybridization with a DNA probe, and infectivity
10 characteristics. The two step test, (in any desired order) however, is a minimum, in order to distinguish over the possibility that the samples may be infected with HBV.

Lack of cross reactivity with polyvalent IgG
15 anti-HBsAg is also indicative of the presence of NANB virus and can be added to the battery of the aforementioned tests. It is however, not conclusive evidence since positive identification such as concentration of antigen is still needed to confirm its
20 presence.

The invention lends itself to the preparation of kits useful in the diagnosis of NANB hepatitis. For example, such a kit may comprise a carrier being compartmentalized to receive one or more container
25 means therein, including a first container containing a monoclonal IgM antibody having immunoreactivity towards said NANB virus; and a second container containing a monoclonal antibody having immunoreactivity towards HBsAg but no immunoreactivity towards the NANB virus.

30 The kit may also comprise a third container means containing detectably labeled HBV-DNA probe,



-21-

and/or additional container means containing another monoclonal antibody having immunoreactivity towards HBsAg but no immunoreactivity towards the NANB virus.

5 Detectably labeled HBV-DNA may also be present in the kit in another container.

10 The use of hybridization techniques initially with purified cloned HBV-DNA can be utilized to clone the DNA of NANB hepatitis viruses with partial sequence homology to HBV-DNA. This is based on the finding that even under very stringent hybridization conditions, the HBV-DNA probe is capable of detecting NANB virus in both human and chimpanzee serum. With purification of the virus by the monoclonal antibody affinity column described herein, the DNA of the virus can be extracted and cloned in bacterial plasmids such as pBR 322 or bacteriophages such as bacteriophage λ .

20 A series of restriction endonucleases are used to cleave the DNA into specific segments with known specific 5' and 3' ends by recognition of specific hexanucleotide sequences in double-stranded DNA. These DNA fragments can then be introduced into plasmids or bacteriophages treated with the same restriction enzymes to produce chimeric recombinant DNA molecules. These recombinant DNA molecules are introduced into E. coli, amplified and produced in large amounts. Recombinants containing NANB virus DNA sequences related to HBV-DNA are identified by molecular hybridization using standard screening procedures. A large group of such clones can then be used to find additional clones with NANB virus sequences only slightly related to HBV-DNA. By this approach, the entire molecular structure of NANB



-22-

hepatitis virus(es) can be reconstructed. With this information and these clones, new recombinant DNA clones can then be prepared which are unique for NANB hepatitis virus(es).

5 The availability of purified isolated NANB virus, substantially free of cellular components and other viral or non-viral components, allows for the preparation of an NANB vaccine. The vaccine can be prepared according to a number of well known methods in
10 the art. Thus, a vaccine can be prepared from the whole live virus or from immunologically active but non-pathogenic subcomponents thereof, such as capsids and the like, obtained by splitting with enzymes or solvents. Chemically attenuated live or killed viral
15 vaccines can also be used, for example, by the treatment of virus with propio lactone, dilute formalin(i.e., conc. less than 1%), ethylene amine, halogenated hydrocarbons, and the like. These agents decrease virus pathogenicity while allowing the
20 material to retain immunogenicity.

Another technique for attenuating the virulence of the virus is to develop an avirulent or slow growing strain, or a mutant incapable of sustained replication in the host. This is generally known in the art as
25 "genetic attenuation", and can be done by genetic manipulations or by serial passage. For example, the production of live attenuated viruses can be carried out by adapting the isolated virus to cultures containing tissue cells and attenuation for example by
30 10-200 passages in such cultures, after which said viruses multiply and a vaccine is then prepared. Another method of producing live vaccine is to select and culture clones. If the infected cells are used for



-23-

the production of the live vaccine, it is advantageous to release the virus from the cells. Techniques for preparing vaccines are generally detailed in a publication such as "Newcastle Disease Vaccines: Their
5 Production and Use", Allan, W.H., J.E. Lancaster and B. Toth; Food and Agricultural Organization, Rome 1978.

The vaccines, whether live or attenuated, in their many different forms, can be prepared in suspension in a manner known per se with a
10 pharmacologically acceptable vaccine carrier, such as a bio-acceptable oil. It is advantageous to add thereto a stabilizer, particularly if a dry preparation is prepared by lyophilization. An adjuvant such as aluminum hydroxide may be added. The stabilizing agent
15 can be a carbohydrate such as sorbitol, mannitol, starch, dextran or glyucose; a protein like albumin or casein; a protein-containing agent like bovine serum or skim milk, and a buffer such as an alkaline metal phosphate. 1-100 µg of virus can normally be present
20 in such composition per unit dosage.

The vaccine can be administered to animals, especially humans, to prevent the same from developing NANB hepatitis. Vaccines (1-100 µg of antigen) may be administered intramuscularly followed by 2nd, 3rd and
25 even more boosts at 2 two month intervals. It should be noted that vaccines may be given subcutaneously or intravenously and the route of administration, dosages, and time between primary immunization and secondary boosts will depend on the immunogenicity and
30 characteristics of the viral antigens employed.

Having now generally described this invention, the same will become better understood by reference to



-24-

certain specific examples which are included herein for purposes of illustration only and are not intended to be limiting of the invention unless otherwise specified.

5

EXAMPLE 1

Monoclonal IgM Radioimmunoassay for Hepatitis B
Surface Antigen: NANB-Binding Activity in Serum
that is Unreactive with Polyvalent Antibodies

MATERIALS AND METHODS

10

Patients. Patient A was a 26-year-old man with acute hepatitis (AH). At the time of study the serum glutamic-oxaloacetic transaminase (SGOT; aspartate aminotransferase) was 2161 international units (IU)/ml (normal < 50), bilirubin was 9.2 mg/100 ml (normal < 1.0), and alkaline phosphatase was 119 IU/liter (normal < 45). His disease resolved over 2 months. Patient B was a 65-year-old man with chronic active hepatitis (CAH). He developed AH 2 months after multiple transfusions for gastrointestinal hemorrhage due to a duodenal ulcer. Liver biopsy showed a histologic pattern consistent with acute viral hepatitis with submassive necrosis. The patient improved, with SGOT, bilirubin, and alkaline phosphatase values returning to normal over several weeks. However, 2 months later he was again icteric and symptomatic; liver biopsy showed CAH with postnecrotic cirrhosis. For the last 4 years his disease has remained active, with SGOT values ranging between 45 and 221 IU/ml, with mildly increased alkaline phosphate levels. Patient C was a 42-year-old woman blood donor. Her physical examination and SGOT, bilirubin, and alkaline phosphatase were normal. Patient D was a 58-year-old man with HBsAg-positive CAH



-25-

proven by liver biopsy. Patient E was a 36-year-old man with AH. The SGOT was 650 IU/ml, bilirubin was 2.4 mg/100 ml, and alkaline phosphatase was 121 IU/liter at the time of study.

5 Patient E had no serologic markers for hepatitis A or B [negative for HBsAg, antibodies to hepatitis B core antigen (anti-HBc), anti-HBs, and IgM antibodies to hepatitis A antigen (anti-HA); tested by Abbott RIAs] during the acute phase of his disease.

10 Patient A was positive for anti-HBc and anti-HBs but negative for HBsAg and IgM anti-HA. Patient B was also negative for HBsAg anti-HBs and IgM anti-HA during AH. However, after the development of CAH he became positive for anti-HBc and anti-HBs but not HBsAg and

15 remained seropositive for these antibodies for the last 4 years in the setting of active liver disease. He was negative for anti-HA IgM. Patient C had no serologic markers for hepatitis A or B. Patient D was positive only for HBsAg and anti-HBc.

20 Patients A, B, C and E were selected for more detailed study because of the high binding activity exhibited by their serum in a 5D3-5D3 monoclonal sandwich RIA. It should be noted that patient B serum was highly positive in the RIA during AH and CAH and he

25 was consistently identified by the assay under code. Patient C was of special interest; her blood was considered to have transmitted acute hepatitis with no serologic markers of hepatitis B or A. Ten units of blood were transfused to the recipient and under code


30 her serum was the only one of the eight units available for study that was reactive in the monoclonal assay. Patient D was selected as a control because his serum was highly reactive for HBsAg with both the monoclonal



RIA and commercial RIA (AUSRIA II, from Abbott).

Affinity Purification. Studies were performed to isolate from serum the high binding activity detected in the 5D3-5D3 monoclonal RIA. Affinity
5 columns of monoclonal 5D3 IgM anti-HBs were prepared by coupling 2-4 mg of IgM per ml of cyanogen bromide-activated Sepharose 6B®. Serum (20-50 ml) from each patient was placed over the columns and incubated for several hours at room temperature; the columns were
10 then extensively washed with phosphate-buffered saline (P_i /NaCl) (pH 7.2). Subsequently, 1- to 2-ml fractions were collected by elution with glycine HCl buffer (pH 2.6). The pH of each fraction was adjusted to 7.4 with 0.1 M NaOH and the binding activity was determined on
15 the eluates by the monoclonal and AUSRIA II RIAs. Peak fractions exhibiting the highest binding activity were pooled and concentrated approximately 100-fold by the Micro-ProDiCon device (Bio-Molecular Dynamics, Beaverton, OR) for further studies as outlined below.

20 Immunoelectron Microscopy. Serum samples (3-5 ml) and 5D3-affinity-purified material from the patients with acute or chronic hepatitis, and serum from normal patients and liver disease controls (individuals with halothane hepatitis, alcoholic
25 hepatitis, or primary biliary cirrhosis who were unreactive in the conventional monoclonal RIA) were incubated for 12 hr at 4°C with 100 µg of 5D3 IgM purified by Sepharose 4B chromatography. The incubation mixture was centrifuged at 12,000 X g for 1
30 hr, the supernatant was decanted, and the precipitate was resuspended in 30 µl of P_i /NaCl. Drops (5-10 µl) were applied to colloidalion/carbon-coated specimen grids, negatively stained with 2% potassium



-27-

phosphotungstate (pH 7.2), and examined with a JEOL 100B electron microscope. Additional controls consisted of serum and 5D3 affinity-purified material incubated with 100 μ l of serum having an anti-HBs titer of 1:500,000 by passive hemagglutination. The latter serum was obtained from a multi-transfused hemophiliac.

Antigenic Characterization. In order to further define the antigenic composition of the 5D3 binding material a series of RIAs employing monoclonal IgG and IgM anti-HBs antibodies were developed. In brief, 5D3 IgM anti-HBs was coupled to a solid-phase support, followed by the addition of serial dilutions of serum samples or 5D3 IgM affinity-purified material and 125 I-labeled IC7 and 5C3 (IgG1 and IgG2a monoclonal anti-HBs), 2F11, 1F8 and 5D3 (IgM monoclonal anti-HBs). The reaction mixture was incubated for 4 hr at 45°C and then the solid-phase support was washed with distilled water. Radioactivity (cpm) bound was determined with a Packard gamma counter. The monoclonal antibodies employed in the RIAs were shown to recognize different determinants as demonstrated by the absence of competitive inhibition in HBsAg binding studies [Wands, Jr. et al, Lancet 1:May, 1982 incorporated by reference]. The binding activity exhibited by the samples in the monoclonal RIAs was also compared to that observed with conventional anti-HBs reagents (AUSRIA II). Finally, the 5D3 affinity-purified material was concentrated approximately 100-fold as described above and retested with the AUSRIA II assay. Under these conditions, the NANB antigen became reactive.

Analysis of Polypeptides. Binding material (20-25 μ l) prepared by affinity chromatography from



-28-

patients was applied to NaDodSO₄/10% polyacrylamide gels (Moriarty et al ibid, 78: 2606 (1981)). Sepharose 4B column-purified 5D3 IgM anti-HBs served as control. Therefore, the polypeptide profiles on the 5 gels of the affinity-purified material derived from patients A, B and C and the HBsAg-positive patient were compared with CAH (patient D).

Molecular Weight Determination. Experiments were performed to determine the approximate molecular weight of the 5D3-binding material. Serum samples (10-15 ml) from patients B and C were placed over Sepharose 4B columns and eluted with P_i/NaCl. The molecular weight markers were blue dextran, IgM, IgG and myoglobin. Aliquots of the fractions were tested in 15 the 5D3-5D3 monoclonal RIA and the binding activity was compared to the elution profiles of the molecular weight markers. The fractions exhibiting the highest binding activity were pooled, concentrated, and immunoprecipitated with 5D3 as noted above and examined 20 by electron microscopy.

RESULTS

Figure 1 depicts a typical binding profile of the various fractions eluted from the 5D3 IgM anti-HBs affinity columns as measured by the 5D3-5D3 monoclonal 25 RIA. Binding activity was recovered from serum after elution with glycine HCl buffer and, as can be seen in Table 1, the amount of radioactivity bound in the peak fractions was higher than that obtained in the unfractionated serum.



Table 1. Antigenic characterization of 5D3 binding activity by polyvalent anti-HBs reagents

Patient	Binding, cpm					
	Serum*		5D3 affinity purified†		Concentrate‡	
	5D3-5D3	AUSRIA II§	5D3-5D3	AUSRIA II§	5D3-5D3	AUSRIA II§
A	15,215	142	34,162	121	72,510	8216 (56)
B	5,610	128	22,300	118	66,721	4432 (42)
C	8,126	142	34,136	102	54,613	2167 (71)
D	22,416	20,618 (11,210)	46,198	26,210 (12,617)	-	-
E	34,259	137	-	735	-	5792 (117)

-29-

- * One hundred microliters of serum tested in the simultaneous 5D3-5D3 monoclonal or AUSRIA II RIA. Results are positive if the cpm bound are greater than 210 or 350, respectively.
- + Binding activity isolated from 30-50 ml of serum by affinity chromatography. In each RIA, 100 µl was tested.
- † Peak binding fractions (see Fig. 1) were pooled (5-7 ml) and concentrated to 50 µl by Micro-ProDiCon. In each RIA, 10 µl was tested.
- § The numbers in parentheses represent the values obtained in AUSRIA II after a 12-hr preincubation with purified 5D3 IgM monoclonal anti-HBs.



-30-

No binding activity was observed with conventional polyvalent anti-HBs reagents. Furthermore, the fractionated serum was devoid of binding activity after passage over the columns and elution with P_i /NaCl as measured by the monoclonal RIA. (FIGURE 1)

Some of the antigenic characteristics of the 5D3-binding material were determined in this Example. (See also below). In one study five monoclonal RIAs were employed as shown in Fig. 2. Fig. 2 Right is a semilogarithmic plot of the binding profile with serial dilutions of serum tested in RIAs using the monoclonal IgM and IgG anti-HBs antibodies (5D3, 2F11, 1F8, 1C7 and 5C3). All immunoassays showed high reactivity in the patient with HBsAg-positive CAH. In contrast, only the 5D3-5D3 RIA identified serum from patients B and C as positive, as shown by the absence of significant binding activity when the four other monoclonal RIAs were used (Fig. 2 Left). These findings indicate that the reactivity of these sera in 5D3-5D3 assay was the result of a specific antigen-antibody interaction and not just due to nonspecific binding of serum to murine monoclonal IgG and IgM anti-HBs.

Additional antigenic properties of the 5D3-binding material are also shown in Table 1. The degree of binding activity increased in the 5D3-5D3 assay as the serum samples from patients A, B and C were affinity purified and further concentrated. It is of interest that when all four specimens were concentrated approximately 100-fold (by volume) they showed strongly positive results with the polyvalent anti-HBs reagents (AUSRIA II). However, this binding activity was blocked by preincubation of these samples with 5D3



-31-

monoclonal anti-HBs. In contrast, only a 50% blockage of known HBsAg binding activity was observed in AUSRIA II after preincubation of affinity-purified HBsAg from patient D with 5D3 IgM anti-HBs.

5 The polypeptide profiles of the affinity-purified material from four patients on NaDodSO₄/polyacrylamide gels were compared, as shown in Fig. 3. Some striking similarities in protein bands were observed when comparisons were made among patients
10 A, B, C, and HBsAg derived from patient D. A major 50,000-dalton protein was found to be common to all specimens, although the HBsAg polypeptide migrated slightly ahead of the other 50,000-dalton proteins from patients A, B, and C. Two other polypeptides in the
15 22,000- to 23,000-dalton range appeared to be common components in all four isolates. More importantly, the polypeptide profiles were identical in samples A, B, and C and, although there were some similarities to the polypeptides of HBsAg, as a group there were distinct
20 differences as well.

Finally, the molecular weight of the binding material was approximately 2×10^6 in patients B and C as determined by Sepharose 4B® chromatography.

DISCUSSION

25 This Example shows a study which was designed to compare directly the properties of the binding material detected only in the monoclonal RIA and not in conventional assays (AUSRIA II). If the binding activity measured with the monoclonal RIA was identical
30 to HBsAg, it would have been expected that the conventional assays should also yield positive results



-32-

in view of the known sensitivity of the monoclonal RIA for HBsAg (100 ± 30 pg/ml). The goal of the present study was to assess the relationship, if any, of HBsAg to 5D3 affinity-purified material derived from patients negative in the serum for HBsAg by conventional RIA with AH or CAH, and from a donor whose blood was implicated in transmitting AH to a recipient.


There is no doubt that the monoclonal 5D3 anti-HBs recognized a determinant on HBsAg as shown by the present study and previous observations (Wands et al, PNAS, 78: 1214-1218 (1981)). HBsAg was isolated from serum by the 5D3 IgM affinity column. The immunoreactivity of the isolate was confirmed by the high binding activity measured both in the 5D3 monoclonal and AUSRIA II assays. In addition, when affinity-purified HBsAg was preincubated with 5D3 IgM anti-HBs and the immunoprecipitate was examined by electron microscopy, typical 22- to 25-nm particles were observed. Clumping of the particles and their "spiculated" or "fuzzy" appearance is consistent with the presence of antibody on the surface. Indeed, this observation provides morphologic evidence of the interaction of the 5D3 monoclonal anti-HBs with a specific determinant(s) on HBsAg. The polypeptide profile on NaDodSO₄/polyacrylamide gels of the affinity-purified HBsAg isolate was consistent with previous reports demonstrating a major 50,000-dalton polypeptide and two smaller proteins (23,000 and 27,000 daltons). Finally, the immunoreactivity of the HBsAg isolate was further established by high binding activity in RIAs using the other four monoclonal IgM and IgG anti-HBs antibodies and conventional (commercially available polyvalent anti-HBs AUSTRIA II).



-33-

Some similarities were observed between HBsAg and the 5D3 immunoreactive material (NANB) isolated from patients A, B, C, and E. First, the binding activity recovered from serum by using 5D3-IgM anti-HBs affinity columns and the radioactivity bound in the eluate as measured by the monoclonal RIA was severalfold higher than that measured in serum. Furthermore, concentration of the eluate followed by retesting in the monoclonal RIA yielded even higher binding values. Second, immunoprecipitation of the affinity-purified material revealed distinct particles by electron microscopy. However, no particles were observed in the isolates after the addition of high-titer anti-HBs. The appearance and the size of the NANB particles was similar but not identical to HBsAg. The density of particles on electron microscopic grids was generally less than that observed with the 5D3-HBsAg immunoprecipitate. It should be noted that, as with the 5D3-HBsAg immunoprecipitate, clumping of particles was observed, which presumably represents the presence of 5D3 antibody on their surface. Finally, as shown in Fig. 3, NaDodSO₄/polyacrylamide gel electrophoresis revealed three polypeptides in the same molecular weight range as previously described for HBsAg.

Although the 5D3 IgM anti-HBs binding material (NANB) shared certain properties with HBsAg, distinct differences were noted. These differences were most evident when the antigenic characteristics of the 5D3 immunoreactive material were examined by using other monoclonal IgG and IgM anti-HBs antibodies as well as conventional reagents. Four other monoclonal IgM and IgG anti-HBs antibodies were unreactive with 5D3 binding material when tested in solid-phase RIAs. In



-34-

contrast, all four antibodies were highly reactive with HBsAg in the same RIAs. In this regard, it is noteworthy that 5D3 was coupled to the solid-phase support and the other ^{125}I -labeled monoclonal IgM and 5 IgG anti-HBs served as the indicator probes. It is likely, therefore, that the 5D3 binding material was bound to the solid-phase support but was not detected with the RIAs, suggesting that those epitopes were absent or were not available in sufficient concentration to be identified by the other monoclonal anti-HBs.

Additional examination of the antigenic characteristics of the 5D3-affinity-purified material was performed after concentration of the eluate 15 (approximately 100-fold by volume). After each concentration all these isolates were reactive in AUSRIA II. More importantly, preincubation with 5D3 anti-HBs blocked the binding by the conventional anti-HBs. One possible interpretation is that the 20 conventionally prepared anti-HBs reagents contain small amounts of an antibody like 5D3 IgM anti-HBs or of an antibody of the IgG class that competes for the same epitope. However, these isolates (Table I) possessed high binding values with conventional anti-HBs in 25 commercial RIAs, which suggests some antigenic crossreactivity of the determinants on HBsAg and the NANB 5D3 binding material. In contrast, preincubation of 5D3 with HBsAg resulted in an approximately 50% reduction in binding activity when retested by AUSRIA 30 II. This result was not unexpected, because other unoccupied determinants would be available for binding by the commercial polyvalent anti-HBs. These findings therefore suggest that 5D3 IgM anti-HBs is directed toward a highly represented epitope on HBsAg, but also



-35-

recognizes an epitope shared with the NANB virus.

It should be emphasized that the patients selected for the present Example were part of a much larger group of individuals whose serum gave a positive reaction with the 5D3-5D3 monoclonal assay but not with the commercial RIA. The patient selection was based primarily on very high binding activity of their sera in the monoclonal RIA. Thus, at present it is not clear whether isolates obtained from other sera with lower levels of binding activity will yield the same properties as described above. However, it is evident that the 5D3-IgM material [NANB] may be affinity purified from serum derived from patients with acute and chronic inflammatory liver diseases and even from (a normal) an individual and shows limited crossreactivity with HBsAg when analyzed by conventional anti-HBs reagents. Furthermore this material is not recognized by four other IgM and IgG monoclonal anti-HBs in RIAs, is similar to HBsAg with respect to three polypeptides on NaDodSO₄/polyacrylamide gels, has a molecular weight of approximately 2×10^6 , and appears as distinct particles by immunoelectron microscopy.

EXAMPLE 2

Demonstration Of Previously

Undetected Hepatitis B Viral Related Determinants
In An Australian Aboriginal Population By
Monoclonal Anti-HBs Antibody Radioimmunoassays

Subjects

Approximately three-quarters of the adults and children of Mornington Island, an Aboriginal settlement off the mainland of Queensland in the Gulf of



-36-

Carpentaria were studied. The population is very stable and there is little interchange with the mainland. The subjects and the Department of Health, Queensland, gave permission for blood samples to be
5 taken. Peripheral-blood samples were drawn into heparinised tubes, and the plasma was separated by centrifugation.

Production and Characterization of Monoclonal
Anti-HBs Antibodies

10 The immunization protocols, characteristics and purity of the immunization antigen (HBsAg), cell-fusion technique, and growth and cloning of hybridomas producing anti-HBs antibodies have been described previously (Wands et al, Gastroenterology, 80:225-
15 232). The anti-HBs antibodies have been characterized with respect to specificity for determinants on HBsAg, ability to agglutinate red blood cells coated with HBsAg (subtypes adw and ayw), antibody class and subclass, and affinity for HBsAg-associated epitopes.
20 Two IgM and two IgG monoclonal anti-HBs antibodies were selected for this study because they recognize all known subtypes of HBsAg and have very high affinity constants for HBsAg determinants and also recognizes NANB antigenic activity: the monoclonal antibodies 5D3
25 and 3D4 (IgM), 5C3 (IgG_{2a}), and 2C6 (IgG₁) have affinity constants of 4×10^{11} , 8×10^{10} , 4×10^{10} , and 2×10^{10} litres/moles per molecule, respectively.

Monoclonal IgM and IgG Anti-HBs Radioimmunoassays (Test Procedures)

30 Previous studies (Shorey, J. et al Hepatology 1:546 (1981) (Abst.)) have established that the 5D3 IgM



-37-

monoclonal anti-HBs antibody recognizes all known HBsAg subtypes and, more importantly, has the highest affinity constant of the anti-HBs antibodies measured. 5D3 anti-HBs was coupled to a solid-phase support, and the other IgM and IgG antibodies were radiolabelled with iodine-125 to a specific activity of 4-10 $\mu\text{Ci}/\mu\text{g}$. Before iodination the antibodies were purified from ascites fluid by staphylococcal-protein-A affinity chromatography for IgG and 'Sephacrose-4B' chromatography for IgM. For the monoclonal radioimmunoassays, approximately 50 ng 5D3-coated beads were incubated with 100 μl serum and 100 μl (150,000 cpm) radiolabeled monoclonal anti-HBs for 16 h. The solid-phase support was washed three times with distilled water, and the radioactivity bound to the bead was measured by a Packard gamma well counter.

All serum samples were evaluated with the 5D3-5D3 "simultaneous sandwich" radioimmunoassay in which the antibody on the solid-phase support and the radiolabeled indicator antibody are the same. This assay design is the most sensitive for detection of an HBsAg-related determinant. Once high binding activity was demonstrated in serum, three other monoclonal radioimmunoassays were performed in which radiolabeled 3D4, 2C6, or 5C3 anti-HBs was the indicator probe. It was possible, therefore, to determine whether there were additional antigenic determinants in the 5D3-immunoreactive material which could be detected by the other high-affinity monoclonal antibodies. All serum samples from the Aboriginal population were also tested for HBsAg, anti-HBs, and hepatitis B core antibody (anti-HBc) by commercial radioimmunoassays ('Ausria II', 'Ausab', and 'Corab', respectively; Abbott Laboratories, North Chicago, Illinois).



Analysis of HBs-Ag-related Determinants

Competitive-inhibition studies were carried out to determine whether the four monoclonal anti-HBs antibodies recognize the same, closely related, or
5 separate antigenic determinants on HBsAg. For these investigations HBsAg (subtypes adw and ayw) was coated to a solid-phase support and was incubated for 16 h with a constant concentration of radiolabeled 5D3 (150,000 cpm) and various amounts of purified unlabeled
10 5D3, 3D4, 2C6, or 5C3 anti-HBs. It would be expected that high concentrations of unlabeled 5D3 would completely inhibit binding of radiolabeled 5D3 to its determinant on HBsAg. If, when another monoclonal anti-HBs such as 2C6 is incubated with radiolabeled
15 5D3, there is no inhibition of 5D3 binding to HBsAg, it may be concluded that 5D3 and 2C6 bind to different determinants. To provide further evidence in support of this conclusion reverse experiments were performed in which, for example, 2C6 was radiolabeled and
20 incubated with various concentrations of unlabeled 5D3. If there is no inhibition of binding of the labeled antibody in the presence of a high concentration of the other, unlabeled antibody, the two antibodies must be directed against distinct and
25 separate determinants on the hepatitis-B-virus related protein.

Results

The IgG anti-HBs antibodies 2C6 and 5C3 had no effect on the binding of 5D3 to an HBsAg-related
30 determinant (Fig. 4), whereas 3D4, an IgM anti-HBs, partially inhibited 5D3 binding. Additional



-39-

experiments confirmed that 5C3, 2C6, and 5D3 recognized distinct and separate determinants on HBsAg. There was some antigenic cross-reactivity between the 5D3 and 3D4 epitopes; 3D4 binding was not, however, influenced by the two IgG anti-HBs antibodies (5C3 and 2C6). The four monoclonal radioimmunoassays used in this study detect three separate epitopes and one partially cross-reactive epitope on HBsAg.

Approximately 50% of the study population had been exposed to HBV as shown by the presence in serum of HBsAg, anti-HBs and anti-HBc, or both antigen and antibodies (Table 2).




TABLE 2 - HEPATITIS B VIRUS MARKERS IN MORNINGTON ISLAND RESIDENCE

		No. with marker (%)			
		Positive by			
Patient		Anti-HBs			
5	group*	Anti-HBs	AntiHBc	anti+HBc	Ausria II 5D3 RIA
	Adult men (n=96)	23(24.9)	7(7.3)	27(28.1)	5(5.2) 7(7.3)
10	Adult women (n=73)	13(17.8)	7(9.6)	15(20.5)	5(6.8) 10(13.7)
	Male children (n=57)	5(8.7)	1(1.7)	15(26.3)	2(3.5) 5(8.8)
15	Female children (n=50)	4(8.0)	1(2.0)	6(12.0)	1(2.0) 6(12.0)
	Total + (n=316)	51(16.1)	22(6.98)	73(23.0)	14(4.4) 31(9.8)

20 * Analysis on 276/316 subjects for whom data on age and sex were available.

+ Total population was 316 subjects.



-41-

14 subjects were positive for HBsAg by the commercial radioimmunoassay (AUSRIA II); all were highly positive by the 5D3-5D3 simultaneous sandwich monoclonal radioimmunoassay, as were 17 other subjects 5 who were negative by commercial RIA (AUSRIA II). There was a higher frequency of hepatitis B VIRUS markers in male subjects of all ages than in female subjects.



TABLE 3- DEMONSTRATION OF HBV-RELATED VIRAL
DETERMINANTS BY MONOCLONAL-ANTIBODY BINDING

Sample No.	Anti- HBs	Anti- HBc	S/N* measured by				
			Ausria II	5D3- 5D3	5D3- 5D3	5D3- 2C6	5D3- 3D4
1	-	-	0.6	8.3	4.1	0.7	6.4
2	-	-	0.4	11.0	1.0	0.5	4.3
3	+	+	0.7	4.1	0.5	4.1	0.6
4	-	-	1.1	4.1	8.0	1.2	6.4
5	+	+	0.3	13.0	0.6	0.3	17.0
6	+	-	0.1	4.0	8.0	0.5	1.3
7	-	-	0.9	23.0	2.1	1.1	3.5
8	+	+	1.3	7.1	0.6	0.7	2.8
9	-	+	0.7	5.4	5.4	0.7	2.0
10	-	-	0.9	9.7	0.3	0.9	6.3
11	+	-	0.8	3.1	0.6	0.5	2.1
12	-	-	1.1	11.0	0.9	0.9	1.9
13	+	+	1.3	15.0	0.6	0.6	2.4
14	-	-	0.4	3.9	0.3	0.3	1.6
15	+	-	0.4	11.0	1.4	0.4	4.1
16	-	+	0.8	5.5	0.4	0.5	2.0
17	+	+	1.7	7.0	0.8	0.4	1.7
Total† (n=17)	8(47%)	7(41%)	0(0%)	17(100%)	5(29%)	1(6%)	12(71%)
Controls†							
AA† (n=14)	0(0%)	14(100%)	14(100%)	14(100%)	14(100%)	14(100%)	14(100%)
BD (n=100)	7(7%)	5(5%)	0	1(1%)	1(1%)	0	2(2%)

* S/N represents signal/noise calculated as means cpm bound in experimental samples divided by the mean cpm of the negative controls. Result considered positive if S/N >2.0.

† No. (%) of subjects positive.

‡ Australian Aborigines' samples positive by AUSRIA II radioimmunoassay. All samples were reactive with monoclonal anti-HBs IgM and IgG antibodies.

BD = blood donors.




-43-

In addition to the 14 HBsAg-positive subjects shown in Table 2, the 5D3-5D3 monoclonal radioimmunoassay demonstrated significant binding activity in 17 other subjects. The results of further monoclonal-antibody analysis of these samples are given in Table 3. Several patterns were observed. For example, positive results were obtained in sample 1 by the 5D3-5D3, 5D3-3D4, and 5D3-5C3 radioimmunoassays but not by the commercial radioimmunoassay (AUSRIA II) or by the 5D3-2C6 radioimmunoassay. Samples from other subjects, such 2, 5, and 8 were HBsAg-positive by only the 5D3-5D3 and 5D3-3D4 assays; in each of these cases 5D3 was coupled to the solid-phase support and the other ^{125}I -labelled monoclonal IgM and IgG anti-HBs served as the indicator probe. It is likely that the 5D3-reactive material was bound to the solid-phase support but was not detected by some of the radioimmunoassays with other monoclonal anti-HBs antibodies because the HBsAg viral determinants they recognise were absent or not available in sufficient concentration. It is not surprising that there was binding activity with 3D4 antibody in 12 of 17 (71%) 5D3-positive samples, since competitive-inhibition studies indicated partial antigenic cross-reactivity between the 5D3 and 3D4 determinants. The positivity rate for all the monoclonal radioimmunoassays was negligible in a low-incidence blood-donor population (Table 3); this finding provides further evidence of the specificity of the monoclonal radioimmunoassays for HBsAg-related determinants.

Discussion

It has been found that more than 50% of the Aboriginal community on Mornington Island had been



-44-

exposed to HBV. A very high rate of infection with HBV would be expected in confined Aboriginal communities such as that on Mornington Island because of the amount of close contact within household group, the poor
5 socioeconomic conditions, and the very high incidence of venereal disease.

The specificity of the high-affinity IgM and IgG monoclonal antibodies has been confirmed by this Example. The antibodies were prepared against HBsAg,
10 and each has been demonstrated to react specifically with known HBsAg subtypes. Competitive-inhibition experiments indicate that the antibodies recognise distinct and separate determinants on HBsAg. All serum samples which reacted with conventional polyvalent
15 anti-HBs antisera (AUSRIA II) also reacted strongly with the four monoclonal anti-HBs IgG and IgM antibodies, and serum samples from a control caucasian population known to have a low incidence of HBV exposure, reacted infrequently with the monoclonal
20 antibodies. The dilution curves for antibody binding to HBsAg in serum are remarkably similar, which indicated that these viral determinants are present in high frequency on HBsAg and that they are distributed homogenously in the population.

25 The 17 subjects whose serum was reactive in the 5D3-5D3 monoclonal radioimmunoassay but was negative when tested by polyvalent conventional anti-HBs antisera are particularly relevant to this invention. The 5D3-5D3 radioimmunoassay has a sensitivity of 100 pg/ml
30 serum for an HBsAg-related determinant, which represents a sensitivity several times greater than that of the commercial radioimmunoassay. Therefore, some of the positive results may be explained on the



basis of the greater sensitivity of the assay but other positive results cannot be explained by this mechanism. Other monoclonal radioimmunoassays using antibodies which recognize different HBsAg determinants have demonstrated enhanced binding activity in substantial numbers of the 5D3-positive samples. These findings add further support to the concept that the 5D3 binding activity is related to the presence of the NANB hepatitis-B-related viral determinants in serum.

10 In Example 1 were investigated serum samples that exhibited high binding activity in the 5D3-5D3 monoclonal radioimmunoassay but were negative by AUSRIA II, and some of the properties of the binding material were characterized, supra. The finding of antigenic

15 determinants recognized only by high-affinity monoclonal antibodies in a high proportion of the Mornington Island population without conventional HBV markers, indicates that there are additional viruses in this community antigenically related to HBV but not

20 previously detected, i.e., NANB virus. This will be proved by subsequent example (See below).

EXAMPLE 3

Demonstration of NANB viral DNA in Human Serum

Materials and Methods

25 Serum Specimens and RIA's - Serum Specimens and RIA's were those of Examples 1 and 2.

Because several individuals who were reactive only in monoclonal RIAs had anti-HBs and anti-HBe antibodies in the serum, additional experiments were

30 performed to ascertain sensitivity of the monoclonal RIA for an HBsAg-related determinant in the HBsAg-anti-



-46-

- HBs immune complexes formed at various antigen/antibody ratios. In these investigations, several chronic carriers of HBsAg were selected, and serial dilutions were made of their serum (with HBsAg-negative serum).
- 5 Binding activity was measured in each specimen by monoclonal RIAs and was compared to that obtained with polyvalent anti-HBs antibodies (AUSRIA II.) Dilution of HBsAg-positive serum (200 μ l) was then incubated with 25 μ l of serum from a multiply transfused
- 10 hemophiliac patient (with an anti-HBs titer of 1-2.2 X 10^6 by passive hemaagglutination) for 12 hr at 20°C. After this incubation, the RIAs were performed; monoclonal anti-HBs and AUSRIA II for HBsAg and AUSAB for anti-HBs levels.
- 15 HBV DNA Hybridization Studies - For molecular hybridization studies, 10 μ l aliquots of human serum were applied to nitrocellulose filter sheets and denatured and fixed to the filter with 0.5 M NaOH. The material was neutralized, on the filter, with 0.5 M.
- 20 Tris-HCL pH 7.4-1.5 M NaCl, digested with proteinase K (200 μ g/ μ l) in 0.3 M NaCl/0.03 M Na citrate, air dried, and baked in vacuo at 80°C for 2 hr. The bound DNA was prehybridized and hybridized with HBV [32 P] DNA. For these experiments, recombinant cloned HBV
- 25 DNA (13,250 base pairs) was repurified from plasmid pAO1 HBV DNA by digestion of the plasmid with restriction endonuclease EcoRI, followed by agarose gel electrophoresis and electroelution of the purified HBV DBA band. HBV DNA was labeled with [32 P]dCTP and
- 30 [32 P]dATP to a specific activity of 2-4 X 10^8 cpm/ μ g of DNA by nick-translation. Hybridization was performed in 0.75 M NaCl/0.075 M Na citrate/0.02% polyvinylpyrrolidone/0.02% Ficoll/0.02% bovine serum albumin containing denatured calf thymus
- 35 DNA (150-200 μ l) and heat-denatured HBV [32]DNA (1 X 10^6




-47-

cpm/ml) at 65% for 24-36 hr. After hybridization, the unreacted solution was discarded, and the nitrocellulose filter was washed, dried, and autoradiographed. For control experiments, the test sample was purified HBV DNA, DNA extracted from the PLC/PRF/5 cell line, which contains 5 or 6 copies of HBV DNA per genome equivalent or DNA isolated from serum Dane particles.

Results

Fig. 5 depicts a serial study on a patient with acute hepatitis B and HBsAg-anti-HBs immune complex disease characterized by arthritis, rash and arthralgias. In this figure, the signal/noise ratio for HBsAg (a measure of specific binding activity) is higher with monoclonal anti-HBs (IgM) than with polyvalent anti-HBs (AUSRIA II). More importantly, the monoclonal RIA for HBsAg remained positive for 13 wk after the polyvalent AUSRIA II RIA had become negative. During this period, anti-HBs was present in the serum, suggesting that the monoclonal RIA may detect HBsAg-related determinant in HBsAg-anti-HBs immune complexes formed in anti-HBs excess and that such determinants are not detectable by polyvalent anti-HBs antisera.

To further explore this possibility, two additional studies were performed in which HBsAg-anti-HBs immune complexes were formed in vitro with serum from a chronic HBsAg carrier by the addition of high titer polyvalent anti HBs antibodies. When polyvalent anti-HBs was added to serum from an HBsAg carrier, the monoclonal RIA (anti HBs IgM) remained positive up to a 10-fold greater dilution than did the AUSRIA II RIA.



-48-

When, in place of IgM monoclonal anti-HBs, studies were carried out with IgG monoclonal anti-HBs 5C3 and 5C11, which recognize distinct and separate determinants on HBsAg, similar results were obtained. These findings indicate that monoclonal anti-HBs RIAs can recognize specific viral epitopes in the immune complexes when HBsAg is no longer detectable by polyvalent anti-HBs antibodies.

To determine whether HBV DNA-related sequences were present in serum samples that were positive for HBsAg by RIAs only with monoclonal anti-HBs antibodies, sera (10 µl aliquots) were applied as spots to a nitrocellulose filter sheet and denatured. The DNA material was fixed, hybridized with recombinant-cloned and repurified HBV [³²P]DNA, washed, and autoradiographed. All experiments were performed under code with two investigators independently interpreting the autoradiograms. A series of control samples either positive or negative for HBsAg by AUSRIA II were correspondingly positive or negative for HBV DNA by hybridization, respectively. In several hundred random or unselected specimens from a clinical laboratory analyzed, there was no instance in which the HBV DNA hybridization test was positive when the AUSRIA II RIA was negative.

In a select group of specimens that were positive for HBsAg by RIA with ¹²⁵I-labeled monoclonal anti-HBs IgM (5D3) but were negative by RIA with ¹²⁵I-labeled polyvalent anti-HBs(AUSRIA II), HBV[³²P]DNA hybridization was performed. Three of seven samples were positive for HBV DNA by molecular hybridization. Unlike random specimens from a clinical laboratory, some of these specimens which contain HBsAg-related



antigenic activity as detected by 5D3 anti-HBs RIA (i.e., NANB virus protein) also contained HBV-DNA-related sequences (i.e., NANB virus DNA) as detected by molecular hybridization with purified HBV-DNA.

- 5 To determine the frequency with which sera negative for HBsAg by AUSRIA II but positive for monoclonal anti-HBsAg were positive also for HBV-related DNA sequences, 36 selected specimens previously characterized by monoclonal RIAs and additional samples
10 were hybridized under code with HBV[³²P]DNA (Table 4).



Table 4 Characteristics of patients whose serum was
reactive by both monoclonal RIAs and HBV DNA Hybridization

No.	Diagnosis	AUSRIA II RIA, Monoclonal RIA,		Anti-HBs	Anti-HBc	HBV-DNA Related Sequences
		cpm bound	cpm bound			
1	Acute hepatitis	93	2,641	+	-	+
2	Chronic active hepatitis	141	7,621	-	-	+
3	Post-transfusion hepatitis*	136	2,193	-	-	+
4	Blood donor	147	1,862	-	-	+
5	Blood donor	141	2,613	-	-	+
6	Blood donor†	96	1,562	-	-	+
7	Blood donor	114	684	-	-	+
8	Aus. abor.	119	1,281	-	-	+
9	Aus. abor.	88	691	+	+	+
10	Aus. abor.	94	663	+	+	+
11	Aus. abor.	110	934	+	-	+
12	Aus. abor.	113	2,600	+	+	+
13	Aus. abor.	138	1,084	+	+	+
	Controls (100)	136 ± 17	56 ± 9			

-50-

Aus. abor., Australian aborigine.

*Recipient of blood from patient 4.

†Incriminated in transmitting post-transfusion hepatitis. Serum was not available for analysis for recipient of blood from patient 6.



-51-

Table 4 lists the results together with clinical information and data from other tests including various RIAs. A total of 13 of 36 samples (36%) of specimens from different individuals positive for HBsAg determinants with monoclonal anti-HBs but negative with polyvalent anti-HBs were positive for HBV DNA sequences by hybridization with recombinant-cloned and repurified HBV DNA. Amongst these individual were three patients with acute or chronic hepatitis, four blood donors (two of whom have been implicated in transmission of hepatitis to recipients of their blood), and six Australian aborigines of the isolated population from Mornington Island where HBV infection is endemic (See Example 3).

15 Discussion

In the present Example, the monoclonal RIA is able to bind to viral epitopes in HBsAg-anti-HBs immune complexes formed in the presence of anti-HBs excess. Possible explanations for this phenomenon are: (i) the high-affinity monoclonal anti-HBs may compete more effectively for their determinant(s) than do naturally occurring anti-HBs or (ii) the antibodies may have access to unoccupied determinants in the presence of polyvalent anti-HBs excess. Thus, polyvalent anti-HBs may contain only a small amount of antibody with immunologic properties of 5D3, 5C3 and 5C11 monoclonal antibodies, and, even though immunogenicity is directed against HBsAg-related determinants, the region of immunologic reactivity with the monoclonal antibodies may extend beyond that present in polyvalent antisera. Such a phenomenon could permit detection of HBsAg in immune complexes by monoclonal RIAs, whereas conventional anti-HBs antibodies would demonstrate no binding activity under conditions of anti-HBs excess.




-52-

Although such activity could explain the detection of HBsAg in the presence of excess anti-HBs (Table 4 cases 1 and 9-13) additional consideration is required concerning the positive binding activity observed in patients negative for HBsAg by AUSRIA II RIA who were also anti-HBs negative (Table 4 cases 2-8). Some of these results may be explained by the increased sensitivity of the monoclonal immunoassays for HBsAg-associated determinants as demonstrated by the present and previous examples. In addition, HBsAg in some patients may be present in immune complexes circulating under conditions of anti-HBs equivalence or excess and, as shown here, would be detectable only by monoclonal RIAs.

In terms of the HBV DNA reacting sequences present in 36% of serum specimens positive for HBsAg by monoclonal RIAs but negative by polyvalent RIAs (AUSRIA II) the results indicate that DNA sequences related to or homologous with HBV DNA are present in these specimens. Aside from these selected cases, hybridization with human serum negative for HBsAg by the AUSRIA II RIA has not been detected thus far. Therefore, the present findings do not represent biologically false-positive results.

The presence of both immunoreactive material and hybridizable DNA sequences provides corroborative evidence for the presence of the NANB virus in a significant proportion of these specimens. Positive results were found for both tests under circumstances in which no HBV markers in serum were detected by commercially available RIA (cases 2-7, Table 4). It should be noted that integrated HBV-DNA has been reported in human hepatocellular carcinoma tissue under



-53-

circumstances in which serum of said patients was negative for HBsAg, anti HBs and anti HBc by commercial Abbott kits (Brechot, C. et al, Hepatology, 1, 499 (abstract 9B), 1981 and Brechot, C. et al Hepatology 2, supplement, 27S-34S, 1982 and herein incorporated by reference).

EXAMPLE 4

Infectivity Studies Of Viral Hepatitis In Chimpanzees: Characterization Of NANB Hepatitis B Virus Agents


10 RIA's, HBV-DNA hybridization and antibody specificity were as described in Examples 1-3, supra.
Infectivity Studies

Two chimpanzees were inoculated with one mililiter of serum derived from an individual who had
15 been incriminated in transmitting "non-A, non-B" hepatitis through blood transfusions. Another chimpanzee was injected with 40 mililiters of a clotting factor concentrate previously shown to transmit "non-A, non-B" hepatitis to recipients. The
20 final chimpanzee was inoculated with one milliliter from another individual suspected to harbor a "non-A, non-B" hepatitis agent. Serial studies were performed and immunoreactivity in serum was measured serially by four monoclonal anti-HBs, RIAs, the presence of HBV-
25 related DNA sequences by molecular hybridization analysis, HBsAg by AUSRIA II RIA, antibodies to hepatitis B core antigen (anti-HBc) and anti-HBs (CORAB and AUSAB respectively; Abbott Laboratories, North Chicago, IL) and on selected samples, IgM antibody to
30 hepatitis A virus (HAVAB; Abbott Laboratories).



Results

Figure 6 depicts the observations in a chimpanzee inoculated with the clotting factor concentrate. This animal had previously recovered from HBV infection and was positive for anti-HBs at the time of inoculation and throughout the study period. This chimpanzee was therefor immune to HBV infection as currently recognized and defined. The first evidence of liver injury was apparent on day 40 with a rise in ALT levels to 70 IU/L (nl <38 IU/L); ALT elevations persisted for approximately 35 days. Immunoreactive antigen appeared briefly in low titer following inoculation of 40 milliliters of clotting factor concentrate and then disappeared from the circulation. On day 64 there was a striking rise in serum IgM anti-HBs binding activity from a baseline of 50 CPM to 4010 CPM (S/N ~ 80, nl < 2.1). Antigenemia was subsequently present in the blood for 56 days although titers fell with resolution of the hepatitis. It is noteworthy that ALT levels reached normal values by day 78 but antigen was still detectable in the blood for an additional 42 days. Most importantly, the rise in ALT levels preceded the development of antigenemia and/or viremia by approximately 30 days and thus gives a more accurate description of the incubation period of 64 days (e.g., 24 days after the first ALT rise). Correlations were then made between the appearance of antigen in the blood and the presence of HBV-related DNA hybridizable sequences. As a control, HBV related nucleic acid sequences were not detected during the incubation period by molecular hybridization analysis. In contrast, there was a striking correlation between the rise in antigen titers and the presence of nucleic acid




-55-

material which hybridized to the HBV-DNA probe suggesting that virions were released into the circulation, Figure 7. With respect to other HBV related epitopes, the 5C3 or 5C11 determinants were not
5 detected in serum by RIAs. This observation indicates that NANB virus is antigenically distinct from HBV. Finally, a RIA which employs polyvalent anti-HBs antibodies (AUSRIA II) was unreactive during the course of infection and anti-HBc and anti-HA antibodies were
10 undetectable.

Figure 8 demonstrates the clinical and virologic course of a second chimpanzee with pre-existing anti-HB inoculated with 1 milliliter of serum carrying a "non-A, non-B" agent. In contrast to Figure
15 6, there was no rise in ALT levels during the observation period. The incubation time was judged to be approximately 190 days. The level of antigenemia as reflected by the peak binding activity of the IgM monoclonal RIA was, however, impressive indeed
20 (S/N \sim 175). The period of antigenemia was prolonged (approximately 65 days), and antigen levels became undetectable by day 260. Similar to the first chimpanzee as shown in Figure 6, HBV-related DNA sequences were undetectable during the incubation
25 period but were present by HBV-DNA hybridization at the peak of monoclonal IgM RIA binding activity. Moreover, other HBV related epitopes were absent as determined by the monoclonal RIAs as well as HBsAg (AUSRIA II), anti-HBc and anti-HA antibodies.

30 Figures 9 and 10 illustrate the clinical and virologic course of the final two chimpanzees inoculated with 1 milliliter (each) of serum derived from another individual who had been incriminated in



-56-

transmitting "non-A, non-B" hepatitis. In Figure 9, evidence of liver injury as demonstrated by ALT elevations was apparent on day 50 and persisted with a relapsing pattern for 140 days. Antigenemia appeared on day 92. Antigen titers were, however, falling by day 130 and reached undetectable levels on day 180. The magnitude of peak binding activity by monoclonal RIA was less ($S/N \sim 10$) than that observed in previous studies. It should be noted that anti-HBs was not present at the time of inoculation or during hepatitis infection and recovery. Similar to the chimpanzee shown in Figure 6, the rise in ALT levels precedes the appearance of antigen in the blood by approximately 50 days. HBV-related DNA hybridizable sequences were not detectable now were other HBV associated epitopes, HBsAg, anti-HBc and anti-HA antibodies.

Figure 10 represents the second chimpanzee inoculated with the same serum. In contrast to the pattern seen in Figure 9, ALT elevations were absent. This was similar to the pattern observed in Figure 8. There were three well defined spikes of antigenemia with the highest values occurring on day 164. HBV-related DNA sequences were not detectable during any of the episodes of antigenemia. This chimpanzee was also negative for HBsAg, other HBV related epitopes, anti-HBc, anti-HBs (before, during and after infection) and anti-HA antibodies.

In the present Example it is shown that the agent(s) identified by the techniques of Examples 1-3 is (are) infectious by infectivity studies of viral hepatitis in chimpanzees.

Thus, it has been possible to reproduce the findings in man (See Example 3, Table 4) in an accepted



-57-

experimental animal model of "non-A, non-B" hepatitis. The major observations in the present Example include: 1) three different inocula injected into 4 animals were infectious; 2) the incubation period, defined as the time from inoculation of infections material to the appearance of virus or viral protein in the blood is longer than previously recognized; 3) ALT elevations may precede the appearance of antigenemia by several weeks; 4) antigenemia may occur in the absence of ALT elevations; a phenomenon identical to that observed in man; 5) the presence of antigen in the blood as measured by the monoclonal IgM anti-HB RIAs correlates well with the appearance of HBV-related DNA like sequences by molecular hybridization analysis; 6) the period of antigenemia and/or viremia may persist for weeks to months and usually disappears with recovery; 7) antigenemia is still detectable in the resolution phase of illness when ALT levels are normal, which is similar to HBV infection in man; 8) several episodes of antigenemia may occur during the course of infection; 9) pre-existing anti-HBs was not protective and thus NANB virus is sufficiently different in antigenic composition than HBV. In support of this concept is the finding that polyvalent anti-HBs antibodies (AUSRIA II) and other monoclonal anti-HBs which recognize different HBsAg associated epitopes were unreactive. Taken together these and the previous examples provide strong evidence that NANB hepatitis agents in many circumstances may be a related but distant or distinct variant of hepatitis B virus.

Having now fully described this invention it will be apparent to those of skill in the art that the same can be performed within a wide and equivalent



-58-

range of methods, tests, compositions, procedures and processes without affecting the spirit or scope of the invention or of any embodiment thereof.



Claims

1. Attenuated or inactivated form of a DNA virus which in unattenuated or active form has the following characteristics:

5 A) Molecular weight greater than 2×10^6 Daltons;

 B) Substantial immunoreactivity towards IgM anti-HBs monoclonal antibody 5D3;

 C) Substantially no immunoreactivity towards
10 an anti-HBsAg monoclonal antibody obtained from cell line ATCC CRL 8018;

 D) Concentration dependent immunoreactivity towards polyclonal IgG anti-HBsAg antibodies, which increases with increased concentration of said DNA
15 virus;

 E) Discrete particulate form when observed by immunoelectron microscopy in the presence of 5D3 IgM antibodies;

 F) The DNA of said virus showing partial
20 homology with DNA from hepatitis B virus; and

 G) Said DNA virus showing, in chimpanzees, infectivity having the characteristics of non A, non B hepatitis.

2. The DNA virus of Claim 1, which also has
25 the following characteristic:



-60-

H) substantial immunoreactivity toward an anti HBsAg monoclonal antibody obtained from cell line ATCC HB-8170.

3. The DNA virus of Claim 1 which in its
5 activated or unattenuated form also has the following characteristic:

I) substantially no immunoreactivity toward an anti-HBsAg monoclonal antibody obtained from cell line ATCC HB-8117_.

10 4. The DNA virus of Claim 1 which in ts activated or unattenuated form also has the following characteristic:

J) A polypeptide profile on sodium dodecyl sulfate polyacrylamide gels, when immunoaffinity
15 purified with 5D3 IgM antibody comprising bands at about 50,000, about 23,000 and about less than 20,000 molecular weight.

5. A process for purifying the DNA virus causative of Non A - Non B Hepatitis or a viral protein
20 derived therefrom, from a sample, which comprises:

contacting said sample, comprising said virus or protein together with impurifying amounts of other viral or non-viral components, with a monoclonal antibody having substantial immunoreactivity with said
25 DNA virus, or viral protein but substantially no immunoreactivity with said other components;

selectively binding said DNA virus or viral protein to said monoclonal antibody; and



-61-

removing said DNA virus or viral protein from said monoclonal antibody.

6. The process of Claim 5 wherein said monoclonal antibody is an IgM monoclonal antibody.

5 7. The process of Claim 5 wherein said antibody is 5D3 or is obtained from cell line ATCC HB-8170, and combinations thereof.

8. The process of any of Claims 5, 6, or 7 wherein said monoclonal antibody is immobilized on an
10 insoluble solid phase.

9. The process of Claims 5, 6, or 7 wherein said sample is animal serum.

10. The process of Claim 9 wherein said animal is a human or a chimpanzee.

15 11. The process of Claim 5, 6, or 7 wherein said sample is the fermentation culture of a microorganism which produces said virus.

12. A method of detecting NANB hepatitis producing virus in a sample, which comprises the steps
20 of:

A) confirming the presence of said virus in said sample, and

B) distinguishing said virus from Hepatitis B virus.

25 13. The method of Claim 12 wherein said step



-62-

A) is carried out by immunoassay.

14. The method of Claim 13 wherein said immunoassay is performed using a monoclonal IgM antibody having immunoreactivity towards said NANB virus.

15. The method of Claim 14 wherein said IgM antibody is 5D3 or is derived from cell line ATCC HB-8170.

16. The method of Claim 15 wherein said IgM antibody is 5D3.

17. The method of Claim 14 wherein said immunoassay method is a sandwich immunoassay.

18. The method of Claim 17 wherein said sandwich immunoassay utilizes a first solid phase bound IgM monoclonal antibody and a second detectably labeled IgM monoclonal antibody.

19. The method of Claim 18 wherein both said monoclonal antibodies are 5D3.

20. The method of Claim 12 wherein said step A) is a DNA hybridization.

21. The method of Claim 20 wherein said hybridization step is carried out between DNA from said NANB virus and a detectably labeled HBV or NANB virus DNA-derived probe.

22. The method of Claim 21 wherein said detectably labeled HBV or NANB DNA probe is labeled



with ^{32}P .

23. The method of Claim 12 wherein said step B) is carried out by immunoassay.

24. The method of Claim 23 wherein said immunoassay is performed using a monoclonal antibody having immunoreactivity towards HBsAg but no immunoreactivity toward NANB virus.

25. The method of Claim 24 wherein said monoclonal antibody is an IgM antibody obtained from cell line ATCC CRL 8018.

26. The method of Claim 24 wherein said monoclonal antibody is an IgG antibody obtained from cell line ATCC HB-8171.

27. The method of Claim 12 wherein said step B) is carried out by polyacrylamide gel/SDS electrophoresis of the NANB virus polypeptides.

28. The method of Claim 12 wherein said step B) is carried out by following the infectivity characteristics in a chimpanzee of said NANB virus.

29. The method of Claim 12 wherein said step A) comprises a sandwich immunoassay using a first solid phase bound IgM monoclonal antibody and a second detectably labeled IgM monoclonal antibody both of which are 5D3, and said step B) comprises an immunoassay utilizing a monoclonal antibody obtained from a cell line selected from the group consisting of ATCC CRL 8018 and ATCC HB-8171.



-64-

30. The method of Claim 12 wherein said sample is serum or blood obtained from an animal.

31. The method of Claim 30 wherein said animal is a human.

5 32. The method of Claim 31 wherein said sample is human blood to be transfused.

33. A kit useful for the detection of NANB hepatitis-producing virus in a sample comprising a carrier being compartmentalized to receive one or more
10 container means therein, including a first container containing a monoclonal IgM antibody having immunoreactivity towards said NANB virus; and

a second container containing a monoclonal antibody having immunoreactivity towards HBsAg but no
15 immunoreactivity towards said NANB virus.

34. The kit of Claim 33 which also comprises a third container means containing detectably labeled HBV or NANB-DNA probe.

35. The kit of any of Claims 33 or 34 which
20 also comprises an additional container means containing another monoclonal antibody having immunoreactivity towards HBsAg but no immunoreactivity towards said NANB virus.

36. The kit of Claim 33 wherein said
25 monoclonal antibody in said first container means is 5D3 or is obtained from cell line selected ATCC HB-8170.



37. The kit of Claim 33 wherein said monoclonal antibody in said second container means is obtained from a cell line selected from the group consisting of ATCC CRL 8018 and ATCC HB-8171.

5 38. The kit of Claim 34 wherein said probe is ^{32}P -labeled HBV or NANB-DNA probe.

39. A vaccine composition which comprises an immunogenically active amount of the virus of Claim 1 together with an immunologically inert carrier.

10 40. The composition of Claim 39 wherein said virus is inactivated.

41. The composition of Claim 39 wherein said virus is attenuated.

15 42. The composition of Claim 39 wherein said virus is alive.




FIG. 1










FIG. 2





3/7

FIG.6



4/7

FIG. 3






FIG. 7




5/7

FIG. 8




6/7

FIG. 9



7/7

FIG. 10



INTERNATIONAL SEARCH REPORT

International Application No PCT/US83/01412

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ¹		
According to International Patent Classification (IPC) or to both National Classification and IPC Int. Cl. 3 A61K 39/29; C12N 7/00, 7/02; G01N 33/54 U.S. Cl. 424/89; 435/235, 239; 436/548; 422/61		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
U.S.	424/89; 435/235, 239; 436/548; 422/61	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁶		
Computer Search of Chemical Abstracts File CA - Hepatitis, Non-A, Non-B 1967-1983		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁵	Citation of Document, ¹⁶ with Indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y, P	N, Chemical Abstracts, Volume 98, No. 23, issued 1983 (Columbus, Ohio, U.S.A.), Goodall et al. "Affinity purification of hepatitis B surface antigen using RF-Hbs-1 monoclonal antibody". See page 461, column 2, Abstract No. 196093y, Protides Biol. Fluids 1982 (Pub 1983), 30, 505-8 (Eng.)	1-32, 34-42
Y	N, Chemical Abstracts, Volume 97, No. 1, issued 1982 (Columbus, Ohio, U.S.A.), Collins et al., "Monoclonal antibodies to murine hepatitis virus-4 (strain JHM) define the viral glycoprotein responsible for attachment and cell-cell fusion". See page 338, column 1, Abstract No. 3277j, Virology 1982, 119(2), 358-71 (Eng.)	1-32, 34-42
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁵ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ¹	Date of Mailing of this International Search Report ²	
2 December 1983	21 DEC 1983	
International Searching Authority ¹	Signature of Authorized Officer ¹⁰	
ISA/US	Blondel Hazil	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
X	N, J. Inf. Dis., Volume 145, No. 3, issued 1982 (USA), Robinson W., "The Enigma of Non-A, Non-B Hepatitis". See pages 397-395.	1-4, 39-42
Y	N, Proc. Natl. Acad. Sci., Volume 79, issued 1982 (USA), Wands et al., "Monoclonal IgM radioimmunoassay for hepatitis B surface antigen: High binding activity in serum that is unreactive with conventional antibodies". See pages 1277-1281.	1-32, 39-42
Y	N, The Lancet, issued May 1, 1982 (USA), Wands et al., "Demonstration of Previously Undetected Hepatitis B Viral Determinants In An Australian Aboriginal Population By Monoclonal Anti-Hbs Antibody Radio-immunoassays". See pages 977-980.	1-32, 39-42
X	US, A, 4,291,020, published 22 September 1981, Tabor et al.	1-4, 39-42
X	US, A, 4,271,145, published 2 June 1981, Wands et al.	1-32, 39-42
X, ^P	US, A, 4,356,164, published 26 October 1982 Tabor et al.	1-4, 39-42

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹⁰

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ¹¹

This International Searching Authority found multiple inventions in this international application as follows:

See attached sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☒ No protest accompanied the payment of additional search fees.

PCT/US83/01412

VI. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

- I. Virus, Class 435, subclass 235, Claim 1-4.
- II. Process of purifying a virus, Class 435, subclass 239, Claims 5-11.
- III. Immunoassay, Class 436, subclass 548, Claims 12-32.
- IV. A Kit, Class 422, subclass 61, Claims 33-38.
- V. Vaccine Composition, Class 424, subclass 89, Claims 39-42.